



2024 Recommended Infant Immunization Schedule

For ages Birth - 23 months

Talk with your child's health care provider if you have questions about any vaccine recommendations for your child.

	Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	19-23 MONTHS	
RSV* Respiratory Syncytial Virus	One Dose Depending on Maternal Vaccination Status*									
HepB Hepatitis B	1st Dose	2nd Dose		3rd Dose						
Hib** Haemophilus Influenzae type b			1st Dose**	2nd Dose**	3rd Dose**	4th Dose**				
DTaP Diphtheria, Tetanus, & Pertussis			1st Dose	2nd Dose	3rd Dose	4th Dose				
RV*** Rotavirus			1st Dose***	2nd Dose***	3rd Dose***					
PCV13, PCV15 Pneumococcal disease			1st Dose	2nd Dose	3rd Dose	4th Dose				
IPV Inactivated poliovirus			1st Dose	2nd Dose	3rd Dose					
MMR Measles, Mumps, & Rubella						1st Dose				
VAR Varicella (Chickenpox)						1st Dose				
HepA † Hepatitis A						1st Dose †	2nd Dose †			
COVID-19 †† Coronavirus disease 2019						1 or More Doses of Updated Vaccine ††				
Flu IIV4 ††† Influenza						1 or 2 Doses Every Year †††				

FOOTNOTES

RSV*

Infants born October-March in the U.S. should receive one dose of Nirsevimab if mother did not receive RSV vaccine during pregnancy, if mother's vaccination status is unknown, or if mother received RSV vaccine less than 14 days prior to delivery.

Hib**

Administering a third dose at age 6 months depends on the brand of Hib vaccine used for previous doses. Age 12-15 months need Hib booster dose.

RV***

Administering a third dose at age 6 months depends on the brand of RV vaccine used for previous doses. If brand of initial vaccine is unknown, default to 3-dose series.

HepA †

Two doses of Hep A vaccine are needed for lasting protection. The two doses should be given between age 12 and 23 months. Doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

COVID-19 ††

Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu IIV4 †††

Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

Additional Information

1. If your child misses a recommended vaccine for their age, talk to your child's doctor as soon as possible to see when the missed dose can be given.
2. If your child has any medical conditions that put them at risk for infection (e.g., sickle cell, HIV infection, cochlear implants) or is traveling outside the United States, talk to your child's doctor about additional vaccines that they may need.